SOLUTIONS TO EXERCISES: TRANSFORMING TOOL #1 (the Addition Property of Equality)

IN-SECTION EXERCISES:

- 1. A 'theorem' is a mathematical result that is both TRUE and IMPORTANT (★ that has been proved).
- 2. The universal set for each variable is \mathbb{R} (the set of real numbers). The phrase 'For all real numbers ...' gives us this information.
- 3. 'a equals b (slight pause) is equivalent to (slight pause) a plus c equals b plus c'; or
- 'a equals b if and only if a plus c equals b plus c'
- 4. In this context, 'at least one' means 'one, or more than one'.
- 5. (a) TRUE. In an addition problem, the grouping of the numbers does not affect the result.
- (b) FALSE. For example, choose x = 1 and y = 2. Then, the sentence '1 2 = 2 1' is false.
- (c) TRUE. You can commute the numbers in a multiplication problem, without changing the result. This fact is formally referred to as the **Commutative Property of Multiplication**.
- (d) FALSE. For example, choose x=1 and y=2. Then, the sentence ' $\frac{1}{2}=\frac{2}{1}$ ' is false.
- 6. (a) Add 4 to both sides. (To undo 'subtract 4', apply the transformation 'add 4'.)
- (b) Subtract 7 from both sides. (To undo 'add 7', apply the transformation 'subtract 7'.)
- (c) Add 7 to both sides. (To undo 'subtract 7', apply the transformation 'add 7'.)
- (d) Subtract 7 from both sides. (To undo 'add 7', apply the transformation 'subtract 7'.)
- (e) Subtract 3 from both sides. (To undo 'add 3', apply the transformation 'subtract 3'.)
- (f) Subtract 4 from both sides. (To undo 'add 4', apply the transformation 'subtract 4'.)

7. (a)
$$(1)$$
 (2) (3) (4) (5)

$$5 = 3 + x$$

$$5 - 3 = 3 + x - 3$$

$$2 = x$$

$$x = 2$$

$$5 = 3 + x$$

$$2 = x$$

$$x = 2$$

$$5 = 3 + x$$

$$2 = x$$

$$x = 2$$

$$5 = 3 + x$$

$$2 = x$$

$$x = 2$$

$$x = 2$$
Subtract 3
$$5 = 3 + x$$

$$2 = x$$


$$x = 2$$

$$x = 2$$
For all real numbers x ,
$$5 = 3 + x$$

$$x = 2$$

- 8. (a)
- Suppose that 'a + c = b + c' is true. Then, a + c and b + c live at the same place on a number line.
- Subtracting c from both numbers moves them both by the same amount.
- Since they started at the same place, they'll end up at the same place. Thus, 'a = b' is also true.


The sketch below assumes that c is a positive number:



(b)

- Suppose that 'a + c = b + c' is false. Then, a + c and b + c live at different places on a number line.
- Subtracting c from both numbers moves them both by the same amount.
- Since they started at different places, they'll end up at different places. Thus, 'a = b' is also false.

The sketch below assumes that c is a positive number:



END-OF-SECTION EXERCISES:

For problems 9—12, only one format is illustrated for each solution.

9.
$$4x = 5 + 3x$$
 CHECK:
$$4x - 3x = 5 + 3x - 3x$$

$$x = 5$$

10.
$$x+4=2x-1$$
 subtract x
$$4=x-1$$

$$5=x$$
 add 1
$$5+4\stackrel{?}{=}2(5)-1$$

$$9\stackrel{?}{=}10-1$$

11.
$$3t - 7 + 2t = 4t + 5$$
 CHECK:
$$5t - 7 = 4t + 5$$

$$t - 7 = 5$$

$$t = 12$$

$$3(12) - 7 + 2(12) \stackrel{?}{=} 4(12) + 5$$

$$36 - 7 + 24 \stackrel{?}{=} 48 + 5$$

$$29 + 24 \stackrel{?}{=} 53$$

$$53 = 53$$

12.
$$1 = 8 - t$$

$$1 + t = 8$$

$$t = 7$$
add t
$$t = 7$$
subtract 1

CHECK:
$$1 \stackrel{?}{=} 8 - 7$$

$$1 = 1$$

 $4(5) \stackrel{?}{=} 5 + 3(5)$ $20 \stackrel{?}{=} 5 + 15$